

## Table of Contents

Declaration.....	i
Acknowledgments.....	ii
Summary.....	iv
Table of Contents.....	vi
List of Tables.....	vii
List of Figures.....	x
List of Appendices.....	xii
Introduction.....	1
Material and methods	
i. Sampling, identification and preservation of fish.....	10
ii. Analytical procedures for allozyme electrophoresis.....	13
iii. Analytical procedures for restriction fragment length polymorphism's (RFLP's) of mtDNA.....	20
Results	
i. Identification with phenotypic keys.....	24
ii. Allozyme electrophoresis.....	24
a. General description of alleles and loci.....	26
b. Estimates of alleles per loci, proportion of variant and polymorphic loci and heterozygosity.....	62
c. Estimates of genetic distances and phylogenetic trees.....	63
d. Inbreeding coefficients.....	77
e. Allozymes and different size classes.....	80
iii. RFLP's of total mtDNA.....	81
a. Restriction fragment morphs and divergence estimates.....	88
Discussion	
i. Review and evaluation of results on allozyme electrophoresis.....	93
ii. Review and evaluation of results on RFLP's of mtDNA.....	105
iii. General Discussion.....	111
References.....	126
Appendices.....	139

## List of Tables

Table 1-	Running conditions for allozyme electrophoresis.....	15
Table 2-	Tissue location of enzymes.....	25
Table 3-	a) Allele frequency distribution of <i>Mugil cephalus</i> and <i>Mugil georgii</i> populations along the east Australian coast and from Sri Lanka.	27
	b) Allele frequency distribution of <i>Liza argentea</i> , <i>Myxus elongatus</i> and <i>Aldrichetta forsteri</i> populations along the east Australian coast.	31
Table 4(1-9)-	Observed and expected genotypic distributions and sample sizes for different populations of <i>Mugil cephalus</i> along the eastern Australian waters and Sri Lanka with sample set and pooled data tested for Hardy Weinberg equilibrium for different loci.	34
Table 4(10)-	Observed and expected genotypic distribution for different allozymes for the population of <i>Mugil georgii</i> from Mooloolaba, with polymorphic loci tested for Hardy Weinberg equilibrium.	43
Table 5-	Observed and expected genotypic distributions and sample sizes for different populations of <i>Liza argentea</i> along the eastern Australian waters with sample set and pooled data tested for Hardy Weinberg equilibrium for different loci.	46
Table 6-	Observed and expected genotypic distributions and sample sizes for different populations of <i>Myxus elongatus</i> along the eastern Australian waters with sample set and pooled data tested for Hardy Weinberg equilibrium for different loci.	49
Table 7-	Observed and expected genotypic distributions and sample sizes for different populations of <i>Aldrichetta forsteri</i> along	52

the eastern Australian waters with sample set and pooled data tested for Hardy Weinberg equilibrium for different loci.

- Table 8(1)- Nei's genetic distance estimated between different mullet.....64  
populations using PHYLIP computer package.
- Table 8(2)- Cavalli-Sforza's Chord distance estimated between different .....65  
mullet populations using PHYLIP computer package.
- Table 8(3)- Reynolds, Weir and Cockerham's genetic distance estimated.....66  
between different mullet populations using PHYLIP computer  
package.
- Table 9- Genetic distance estimated between *Mugil cephalus* .....67  
populations from Burrum Heads and Narrabeen  
using 24 loci compared with 19 loci.
- Table 10- (1) Estimates of  $F_{st}$  on polymorphic and variant loci of *Mugil*.....78  
*cephalus* populations.  
(2) Estimates of  $F_{st}$  on polymorphic and variant loci of *Liza*.....79  
*argentea* populations.  
(3) Estimates of  $F_{st}$  on polymorphic and variant loci of *Myxus*.....79  
*elongatus* populations.  
(4) Estimates of  $F_{st}$  on polymorphic and variant loci of .....79  
*Aldrichetta forsteri* populations.
- Table 11- Restriction fragment lengths (bp) identified from different .....82  
*Mugil cephalus* populations from Sri Lanka and eastern  
Australia together with some restriction fragment lengths (bp)  
for *Mugil georgii* from Mooloolaba in eastern Australia.
- Table 12- Restriction phenotypes of 3 mtDNA clones of *Mugil*.....85  
*cephalus* from Sri Lanka and Australia and 2 clones of  
*Mugil georgii* from Mooloolaba in eastern Australia.
- Table 13- A summary of the analysis of inter and intra population .....87  
variation of restriction fragment data of *Mugil cephalus*

populations from eastern Australia and Sri Lanka.

- Table 14- Allele distribution in each species of mullet for the different .....95  
loci studied.
- Table 15- Within species, percent sequence divergence estimates.....112  
based on data derived from restriction endonuclease analysis  
of mitochondrial DNA.
- Table 16- A comparison of sequence divergence estimates derived.....113  
from three mitochondrial methods.
- Table 17- Within genus, mean percent inter specific sequence .....114  
divergence estimates based on data derived from  
restriction endonuclease analysis of mitochondrial DNA..

## List of Figures

- Fig. 1(a)- A sketch of the sampling locations in the Indian and Pacific Oceans.....11
- Fig.1(b)- A detailed map showing the sampling locations from Australia.....12
- Fig. 2(1)-The phylogenetic tree derived through Fitch- Margoliash method with contemporary tips on genetic distances estimated from Nei (1972) on different mullet populations.....69
- Fig. 2(2)-The phylogenetic tree derived through Fitch-Margoliash method on genetic distances estimated from Nei (1972) on different mullet populations.....70
- Fig. 2(3)-The phylogenetic tree derived through Fitch- Margoliash method with contemporary tips on genetic distances estimated from Reynold, Weir and Cockerham (1983) on different mullet populations.....71
- Fig. 2(4)-The phylogenetic tree derived through Fitch- Margoliash method on genetic distances estimated from Reynold, Weir and Cockerham (1983) on different mullet populations.....72
- Fig. 2(5)-The phylogenetic tree derived through Fitch- Margoliash method with contemporary tips on genetic distances estimated from Cavalli-Sforza (1967) on different mullet populations.....73

Fig. 2(6)-The phylogenetic tree derived through Fitch- Margoliash method on genetic distances estimated from Cavalli-Sforza (1967) on different mullet populations.....74

Fig. 3- Restriction fragments produced by Rsa1 digest on western Australian, eastern Australian and Sri Lankan samples of *Mugil cephalus*.....86

Fig. 4- Restriction fragments produced by EcoR1 digest on eastern Australian and Sri Lankan samples of *Mugil cephalus*.....89

Fig. 5- Restriction fragments produced by Sst1 digest on eastern Australian and Sri Lankan samples of *Mugil cephalus*.....90

Fig. 6- Map showing some current patterns of the Indian Ocean. (From Brown *et al* 1989).....109

## List of Appendices

Appendix 1 - CTAB mini preparation for isolating DNA from tissue.....	139
Appendix 2 - Buffers and Stock solutions.....	140
Appendix 3 - Techniques for digestion, electrophoresis and blotting ..... of DNA.	143
Appendix 4 - Techniques for hybridisation .....	146
Appendix 5 - Protocol for extraction of mtDNA from liver of..... <i>Mugil cephalus</i> .	148
Appendix 6 - Recipes for probe labelling.....	150
Appendix 7 - Procedures for washing and exposing membranes.....	152
Appendix 8 - Procedure for developing films.....	154
Appendix 9 - Divergence times estimated on dendogram derived from..... Nei's (1972) genetic distance and KITSCH option on PHYLIP (1991) software package.	155
Appendix 10- Dendogram drawn on Nei's (1972) genetic distance.....	156
Appendix 11- Divergence times estimated on dendogram derived from..... Reynolds <i>et al</i> (1983) genetic distance and KITSCH option on PHYLIP (1991) software package.	157
Appendix 12- Dendogram drawn on Reynolds <i>et al</i> (1983) genetic..... distance.	158
Appendix 13- Divergence times estimated on dendogram derived from ..... Cavalli-Sforza <i>et al</i> (1967) genetic distance and KITSCH option on PHYLIP (1991) software package.	159
Appendix 14- Dendogram drawn on Cavalli-Sforza and Edwards..... (1967) genetic distance.	160