

Contents	Page
ACKNOWLEDGEMENTS	I
LIST OF FIGURES	IV
LIST OF PLATES	VII
LIST OF TABLES	VIII
ABSTRACT	1
Chapter 1. General Introduction	4
1.1 Global overview of the fisheries	4
1.2 Resource depletion	5
1.3 Inland Fisheries of Sri Lanka	9
1.3.1 Reservoir resources and their fisheries	9
1.3.2 Characteristics of reservoir fisheries	10
1.3.3 Fish fauna in reservoirs of Sri Lanka	14
1.4 Unexploited indigenous fish species	19
1.5 Objectives of the present study	26
1.6 Organization of chapters	27
Chapter 2. General Materials and Methods	28
2.1. Study area	28
2.1.1 Land-use patterns and catchment characteristics	36
2.1.2 Seasonal hydrology and draw-down	37
2.2. Sampling	39
2.2.1 Experimental gillnet fishing	39
2.2.2 Shore seining	41

Chapter 3	Page
Exploitation of Small Indigenous Fish Species Using Shore Seines in Three Reservoirs of Sri Lanka	45
3.1. Introduction	45
3.2. Materials and Methods	46
3.3. Results	47
3.4. Discussion	56
Chapter 4	
Population Dynamics of <i>Hyporhamphus limbatus</i> (Belontiiformes, Hemiramphidae) in Two Lowland Reservoirs of Sri Lanka	59
4.1. Introduction	59
4.2. Materials and Methods	60
4.3. Results	63
4.4. Discussion	73
Chapter 5	
Distribution and Abundance of Unexploited Fish Species in Three Sri Lankan Reservoirs	76
5.1. Introduction	76
5.2. Materials and Methods	80
5.2.1. Study area	80
5.2.2. Experimental fishing	81
5.2.3. Water level data Experimental fishing	81
5.2.4. Data analysis	81
5.3. Results	82

	Page
5.4. Discussion	94
Chapter 6	96
Gillnet Selectivity of Small Cyprinids in Three Sri Lankan Reservoirs	96
6.1. Introduction	96
6.2. Materials and Methods	98
6.3. Results	110
6.4. Discussion	
Chapter 7	
Population Dynamics of Non-exploited Fish Species in Three Sri Lankan Reservoirs	113
7.1. Introduction	113
7.2. Materials and Methods	114
7.3. Results	116
7.4. Discussion	133
Chapter 8. General Discussion	135
References	144
APPENDICES	170

LIST OF FIGURES

	Page
Figure 1.1: Status of certain major fish stocks in the world.	6
Figure 2.1: Maps of three reservoirs studied and their major inflows.	31
Figure 2.2: Schematic diagram of the river basin fed to three reservoirs studied in Sri Lanka.	32
Figure 2.3: The hypsographic curves of the three reservoirs.	38
Figure 3.1: Relative importance of the fish species caught in shore seine nets in Minneriya reservoir.	53
Figure 3.2: Relative importance of the fish species caught in shore seine nets in Udawalawe reservoir.	54
Figure 3.3: Relative importance of the fish species caught in shore seine nets in Victoria reservoir.	55
Figure 4.1: The length frequency data of <i>H. limbatus</i> caught in shore seine nets in two reservoirs of Sri Lanka	65
Figure 4.2: The growth curve, length-converted catch curve, selection curve and recruitment pattern of <i>H. limbatus</i> in Minneriya reservoir	67
Figure 4.3: The growth curve, length-converted catch curve, selection curve and recruitment pattern of <i>H. limbatus</i> in Udawalawe reservoir	69
Figure 4.4: The plot of $\ln L_{\infty}$ against $\ln K$ in 95 populations of hemiramphids, and the two populations of <i>H. limbatus</i> in the present study.	70
Figure 4.5: Y'/R curves and the Y'/R isopleths of <i>H. limbatus</i> for Minneriya and Udawalawe reservoirs.	72
Figure 5.1: Relative importance of the fish species caught in gill nets in three reservoirs.	88

	Page
Figure 5.2: Percentage IRI of different species during the sampling dates in inshore and offshore areas in the three reservoirs.	89
Figure 5.3: Bray-Curtis similarity of the species composition in during different sampling dates in the three reservoirs.	92
Figure 5.4: MDS ordination of species composition in the gillnet catches during different sampling dates in the three reservoirs.	93
Figure 6.1: The relationships of Ln catch ratios of fish species to mid-points of length classes.	103
Figure 6.2: Gillnet selection curves of fish species in the three reservoirs studied.	105
Figure 6.3: The relationships between optimal lengths to the gillnet mesh sizes of fish species caught in the three reservoirs.	106
Figure 7.1: Von Bertalanffy growth curves of <i>Amblypharyngodon melettinus</i> studied in 3 reservoirs.	119
Figure 7.2: Von Bertalanffy growth curves of <i>Puntius chola</i> studied in 3 reservoirs.	120
Figure 7.3: Von Bertalanffy growth curves of <i>P. filamentosus</i> studied in 3 reservoirs.	121
Figure 7.4: The length-converted catch curves of <i>A. melettinus</i> studied in 3 reservoirs.	122
Figure 7.5: The length-converted catch curves of <i>P. chola</i> . studied in 3reservoirs	123

	Page
Figure 7.6: The length-converted catch curves of <i>P. filamentosus</i> studied in 3 reservoirs.	124
Figure 7.7: Relative yield-per-recruit as a function of exploitation rate for <i>A. melettinus</i> in 3 reservoirs.	127
Figure 7.8: Relative yield-per-recruit as a function of exploitation rate for <i>P. chola</i> in 3 reservoirs.	128
Figure 7.9: Relative yield-per-recruit as a function of exploitation rate for <i>P. filamentosus</i> in 3 reservoirs.	129
Figure 7.10: The length frequency distributions of <i>A. melettinus</i> studied in three reservoirs.	130
Figure 7.11: The length frequency distributions of <i>P. chola</i> studied in three reservoirs.	131
Figure 7.12: The length frequency distributions of <i>P. filamentosus</i> in studied three reservoirs.	132

LIST OF PLATES

	Page
Plate 1.1: Fiberglass outrigger canoe in Udawalawe reservoir.	12
Plate 1.2: Exotic cichlids, <i>Oreochromis mossambicus</i> and <i>O. niloticus</i> dominate the commercial catch, Udawalawe reservoir.	12
Plate 1.3: Subsidiary fishery for minor cyprinids in Minneriya reservoir.	13
Plate 1.4: Subsidiary fishery for minor cyprinids in Rajanganaya reservoir.	13
Plate 2.1: Satellite image of Sri Lanka showing locations of the three reservoirs.	30
Plate 2.2: Satellite image of Minneriya reservoir.	33
Plate 2.3: Satellite image of Udawalawe reservoir.	33
Plate 2.4: Satellite image of Victoria reservoir.	34
Plate 2.5: Catches of experimental fishing for minor cyprinids in Victoria reservoir.	42
Plate 2.6: Shore seine catch of halfbeak in Udawalawe reservoir.	42
Plate 2.7: Drag net operation for half beak in Udawalawe reservoir.	43
Plate 2.8: Half beaks at a fish stall in Udawalawe together with carps, tilapias and other cyprinids.	43
Plate 2.9: A flock of cormorants in Udawalawe reservoir.	44

LIST OF TABLES

	Page
Table 1.1: List of fish species reported from the reservoirs of Sri Lanka.	15
Table 1.2: The potential yield estimates for minor cyprinids in the different reservoirs.	21
Table 1.3: Past studies on various aspects of small cyprinids in Sri Lankan reservoirs.	22
Table 2.1: Some morphometric, hydrological, physico-chemical and biological parameters of the three reservoirs studies.	35
Table 2.2: Land-uses of the catchments of Minneriya, Udawalawe and Victoria reservoirs.	36
Table 2.3: Hanging ratio, length, depth and area of net panel of gillnets of various mesh sizes used.	40
Table 2.4: Length and depths of three shore seine nets used.	41
Table 3.1: Fish species caught in experimental fishing trials with gillnets and shore seine nets in three reservoirs.	49
Table 3.2: Percentage index of relative importance (% IRI) of fish species caught in shore seines in three reservoirs.	51
Table 4.1 The von Bertalanffy growth parameters of <i>H. limbatus</i> in Minneriya and Udawalawe reservoirs.	64
Table 4.2. The growth performance index (ϕ') of <i>H. limbatus</i> in Minneriya and Udawalawe reservoirs. .	66
Table 4.3. Total mortality (Z), natural mortality (M), fishing mortality (F), exploitation rates and mean sizes at first capture (L_c) of <i>H. limbatus</i> in Minneriya and Udawalawe reservoirs.	68

	Page
Table 5.1: Fish composition expressed as mean catch per unit effort in three reservoirs based on experimental gillnet fishing.	83
Table 5.2: Percentage index of relative importance (%IRI) of fish species caught in gillnets in the three reservoirs.	86
Table 6.1: Catch per unit effort values of minor cyprinids, cichlids and other species caught in different mesh gillnets in the three reservoirs.	99
Table 6.2: Catch per unit effort values of different species in gillnets of different mesh sizes in the three reservoirs.	101
Table 6.3: The regression relationships of $\ln C_2/C_1$ (Y) against mid-length (L in mm) and size ranges of fish.	107
Table 6.4: The estimated optimal lengths, selection factors (SF), standard deviations of selection curves (SD) and corresponding selection ranges of the cyprinid in three reservoirs.	110
Table 6.5: The regression relationships of optimal length (mm) against mesh size (mm) of gillnet for <i>A. melettinus</i> , <i>P. chola</i> and <i>P. filamentosus</i> .	123
Table 7.1 The growth and mortality parameters of <i>A. melettinus</i> , <i>P. chola</i> and <i>P. filamentosus</i> in the three reservoirs studied and in several other Sri Lankan reservoirs.	113
Table 7.2: The mean lengths at first capture (L_c) and exploitation rates (E_{opt}) corresponding to optimal relative yield-per-recruit values and for the three cyprinid species studied in three reservoirs.	122