

# The Nature of the Aerobic Gastrointestinal Bacteria of Cichlid Fish *Sarotherodon mossambicus* (Peters) and *Tilapia Nilotica* (Linnaeus) Grown under Captivity

By

S. K. HAROLD SILVA\* and S. WIDANAPATHIRANA\*\*

## ABSTRACT

Bacteriological examination of the gastrointestinal microflora of two fresh water cichlid fish species (*Sarotherodon mossambicus* and *Tilapia nilotica*) was performed, resulting in the bacteria enumeration of total viable counts of  $1.06 \times 10^7$  /g and  $7.75 \times 10^7$  /g of gastro - bacteria intestinal tract plus contents (wet weight) respectively, by aerobic incubation at  $30 \pm 1^\circ\text{C}$ .

The majority (78%) of the total gut isolates from both fish species was Gram positive mesophilic which is characteristic of the higher ambient temperature in the tropics. These isolates were fastidious in their nutritional requirements and together with the rest are isogenous to bacteria autochthonous to soil and water. The occurrence of such organisms is attributed to the feeding habits of these fish. The gastrointestinal bacteria isolated in this study are transient residents but not 'indigenous' in these cichlid fish.

## Introduction

The gastrointestinal tract of animals has become a popular research area to the ecologist and microbiologist alike. In most animals it is populated by what is termed a 'normal' or 'indigenous' microbial flora. The early concept of the sterility of fish gut (Blake, 1935; Margolis, 1953) has now been replaced by the popular view that feeding fish always harbour viable bacteria in their gastrointestinal tract (Shewan and Hobbs, 1967) and that only migrating fish which undergo fasting may frequently have an empty intestinal tract or at the most a low microbial count (Bramsnaes, 1965). However the presence of a 'normal' bacterial flora in the gut of fish is a controversial issue as investigators believe that fish do not have any natural bacterial flora in their gut, and what is observed has originated from their environment (Wood, 1967) and is a function of the food ingested (Liston, 1956).

The study of gastrointestinal bacteria of fish has enabled investigators to understand biological phenomena underlying spoilage of fish, microbial relationships with the host, bacterial diseases of cultivated fish and food intoxications implicated with consumption of fish. Further, gut bacteria of fish have also served as indicators of faecal pollution of waters inhabited by fish and therefore helped monitoring sanitary conditions of waters.

*Sarotherodon mossambicus* and *Tilapia nilotica* are two fresh water fish species of exotic origin, which were introduced into Sri Lanka as food fish. Of the two the former is reported to constitute a major fishery in itself in the North Central Province and other areas (Mendis and Fernando, 1962), and the latter probably would demonstrate locally the high potential it has exhibited in its native waters (Fryer and Iles, 1972).

\* Microbiology Section, Laboratory Services Division, Sri Lanka Standards Institutions No, 5 Galle Road, COLOMBO 6, SRI LANKA.

\*\* Dept. of Botany, University of Kelaniya, Kelaniya, SRI LANKA.



It was felt that microbiological information of the above two economically important fresh water fish would be most opportune to the developing inland fishery industry of Sri Lanka. Studies on the microbiology of fresh water fish are notably scarce, specially from the tropical region. The present report is the outcome of an investigation on the nature of aerobic gastrointestinal bacteria of the above two fish species grown under captivity.

## Materials and Methods

### *Selection of fish*

Young fish (6–8 months old) of the species *Sarotherodon mossambicus* and *Tilapia nilotica* grown under known culture conditions in domesticated mud ponds were obtained from fresh water fisheries station, Panapitiya, Kalutara. The selected fish which had been caught the previous day and kept overnight in a recovery pond were transported in oxygenated polythene bags containing 1/3 full of clean fresh water. They were in transit for about 1½ hrs and were in very good condition on arrival. The fish were acclimatized and transferred into aquarium tanks containing seasoned fresh water (pH 6.5 – 7.0; temperature 30±1°C). The fish were not fed and the water was neither aerated nor replenished during the course of study. The weight of *S. mossambicus* and *T. nilotica* specimen examined ranged from 32.6g to 93.2 g (average weight 62.9g) and 38.8g to 75.3g (average weight 50.9g) respectively.

### *Post mortem and Bacteriological Examination*

The fish were caught by dip net and killed by delivering a blow to the head (Trust *et al.*, 1979); washed clean. After surface sterilization (Gibbons and Reed, 1930) the gastrointestinal tract in its entirety was removed by adopting aseptic surgery and weighed. It was homogenized in a blender with a standard volume of ¼ strength Ringer's solution.

The resultant homogenate was serially diluted using the same diluent and suitable dilutions are plated on Nutrient agar (Difco) pH 6.8. The seeded plates were incubated aerobically at 30 ± 1°C for 24 to 48 hrs. The total viable counts per gram of intestinal tract plus contents were determined for each specimen.

### *Isolation of bacteria*

Representatives of all the recognisable morphologically different bacterial colonies were picked off from suitable plates. They were purified and the selected pure cultures were transferred into Nutrient agar slants in Bijou bottles. After incubation at 30 ± 1°C for upto 48 hrs these were stored at 4°C as stock cultures.

### *Characterization and Identification of bacteria*

During characterization the preserved stock cultures were recovered in Nutrient agar plates. The morphological characteristics such as pigmentation, staining, shape and arrangement of cells, motility, presence of capsules and endospores of each culture are studied.

The biochemical properties examined included oxidation and fermentation of glucose in Hugh and Leifson's (1953) normal and modified semi solid media; fermentation of and production of acid from glucose, sucrose, maltose, lactose, galactose, salicin, arabinose, mannitol, inositol, starch, xylose and glycerol in broth media; production of acid from glucose, xylose, arabinose and mannitol in modified slants; presence of cytochrome 'C' oxidase, catalase, phosphatase, coagulase, arginine, ornithine and lysine decarboxylases; reduction of nitrate and nitrite; production of indole from tryptophan, H<sub>2</sub>S from Kligler's Iron agar, ammonia from urea and production of acetyl methyl carbinol; detection of mixed acid fermentation; utilization of citrate, malate, aspartic acid, acetate and histidine



as the sole carbon source and reaction to litmus milk. Amylase activity was tested in Nutrient agar having 1% (w/v) soluble starch. Proteolytic activity was tested in Casein agar 1% (w/v) and Gelatine agar 0.4% (w/v) and lipolytic activity was tested in Tween agar having 1% (v/v) Tween 80 (Atlas). Growth in 5% (w/v) NaCl, 10% (w/v) NaCl without added NaCl, growth at pH 5 were tested in Nutrient broth, and growth at 45°C, 65°C were tested on Nutrient agar. Sensitivity to five antimicrobial agents was also tested as necessitated. All the tests were carried out in accordance with the materials and methods given in Harrigan and McCance (1976) and Buchanan and Gibbons (1974).

For the identification of Gram positive coccoid bacteria, Sub Committee Report (1965), Baird-Parker (1962, 1963, 1965 and 1966) were employed. Gram positive endospore forming bacteria were identified on the basis of Gorden and Smith (1949), Smith, Gorden and Clark (1952) and Wolf and Barker (1968). Identification of Gram negative rod shaped bacteria was based on Bain and Shewan (1968), Hendrie and Shewan (1966) and Shewan, Hobbs and Hodgkiss (1960 a & b).

Identification schemes of Cowan (1977) and Buchanan and Gibbons (1974) were adopted for the final identification of the gut isolates.

## Results

The total and average viable counts obtained for the two fish species are illustrated in Table I. The average viable counts recorded for *Sarotherodon mossambicus* and *Tilapia nilotica* were  $1.06 \times 10^7$  bacteria/g (range  $6.11 \times 10^4$  to  $5.76 \times 10^7$  bacteria/g) and  $7.75 \times 10^7$  bacteria/g (range  $1.25 \times 10^6$  to  $3.09 \times 10^8$  bacteria/g) respectively.

TABLE 1

THE TOTAL VIABLE COUNTS OF BACTERIA ISOLATED FROM THE GASTROINTESTINAL TRACT OF *Sarotherodon mossambicus* AND *Tilapia nilotica* (after-h1 24 of incubation at  $30 \pm 1^\circ\text{C}$ )

Sample	Fish	No. of samples	Number of viable bacteria g <sup>-1</sup> (wet weight)	
			Average	Range
Total tract plus contents	<i>S. mossambicus</i>	13	$1.06 \times 10^7$	$6.11 \times 10^4$
				$8.12 \times 10^5$
				$1.43 \times 10^6$
				$1.48 \times 10^6$
				$2.56 \times 10^6$
				$3.38 \times 10^6$
				$3.65 \times 10^6$
				$4.13 \times 10^6$
				$9.50 \times 10^6$
				$1.03 \times 10^7$
Total tract plus contents	<i>T. nilotica</i>	07	$7.75 \times 10^7$	$1.25 \times 10^6$
				$3.48 \times 10^6$
				$8.28 \times 10^6$
				$5.59 \times 10^7$
				$7.56 \times 10^7$
				$8.90 \times 10^7$
				$3.09 \times 10^8$

(Isolation medium - Nutrient Agar pH 6.8)



TABLE 4  
SUMMARY OF BIOCHEMICAL CHARACTERISTICS OF THE BACTERIA ISOLATED FROM  
GASTROINTESTINAL TRACT OF *Sarotherodon mosambicus* AND *Tilapia nilotica*

Bio-chemical characteristics	Number † Positive	Percentage Positive
Pigmentation .. .. .	60(134)	44.77
Catalase .. .. .	132(134)	98.50
Kovacs's Oxidase (Cytochrome 'C') .. .. .	59(134)	44.02
Utilization of Citrate as sole C source .. .. .	56(134)	41.79
Reduction of Nitrate (Nitrate reductase) .. .. .	90(134)	65.69
Hydrolysis of Gelatin (Gelatinase) .. .. .	106(134)	80.91
Hydrolysis of Casein (Casease) .. .. .	93(118)	78.81
Hydrolysis of Starch (Amylase) .. .. .	46(125)	36.81
Hydrolysis of Urea (Urease) .. .. .	72(134)	53.73
Production of Indole (Tryptophanase) .. .. .	9 (134)	6.72
Production of Acetyl Methyl Carbinol .. .. .	52(134)	38.80
Mixed acid Fermentation .. .. .	27(134)	20.14
Oxidative, Hugh & Leifson's test .. .. .	21(134)	15.67
Fermentative Hugh & Leifson's test .. .. .	44(134)	32.84
No reaction, Hugh & Leifson's test .. .. .	69(134)	51.48
Acid from Litmus Milk .. .. .	22(33)	66.66
Production of Arginins Dihydrolase .. .. .	29(33)	87.88
Acid from * Glucose .. .. .	70(134)	52.23
Inositol .. .. .	16(60)	26.66
Mannitol .. .. .	34(87)	39.08
Lactose .. .. .	19(84)	22.62
Acid From ** Glucose .. .. .	32(38)	84.21
Arabinose .. .. .	21(35)	60.0
Mannitol .. .. .	28(35)	80.0
Xylose .. .. .	23(36)	63.83

† Number of strains tested given in parenthesis.

\* Peptone water broth.

\* Modified slant medium (tested for *Bacillus* spp. only)

Further the proteolytic activity shown by a very high percentage of the gut isolates (81% hydrolysed gelatin, 79% hydrolysed casein) and the ability to split urea shown by 54% of the isolates are all characteristics of terrestrial and aquatic bacteria. This evidence strongly suggests that most of the organisms if not all, isolated in this study are similar to those found in soil and water. This is further strengthened by the fact that 52% of the isolates degraded glucose, which is again a characteristic of terrestrial bacteria.

Thus it appears that most of the aerobic organisms isolated from the gut of these fish could have originated in the environment and are probably transient residents in the gut. Obviously, these organisms do not warrant to be placed in the status 'indigenous' organisms of the gut of these fish.

Organisms belonging in the Enterobacteriaceae (*Enterobacter*, *Klebsiella* and *Citrobacter* collectively 8.9%) were encountered, which cannot be considered as indicator organisms in tropical waters (Katugampola and Assim, 1958). The absence of 'faecal' coliform organisms rule out the possible existence of enteropathogenic bacteria among these gut isolates, but it is inconclusive since fish are known to purify themselves when placed in fresh water.



This study has shown that fresh water cichlid fish harbour significant bacterial populations in their gastrointestinal tract. Aerobic and facultatively anaerobic bacteria abound the gut flora which is predominantly Gram positive and mesophilic in character and these organisms reflect the environment of the fish. There is no evidence to indicate that this microflora is 'indigenous' or 'normal' to the gut of these cichlid fish. These organisms are probably transient residents.

## REFERENCES

- BAIN, N. and SHEWAN, J. M. (1968)  
Identification of *Aeromonas*, *Vibrio* and related organisms, in 'Identification methods for microbiologists, Part B' (B. M. Gibbs and D. A. Shapton Eds.), Academic Press, London and New York, 79-84 pp.
- BAIRD-PARKER, A.C. (1962)  
The occurrence and enumeration according to a new classification, of Micrococci and Staphylococci in Bacon and on Human and Pig skin, *J. Appl. Bact.*, 25 (3), 352-361 pp.
- BAIRD PARKER, A. C. (1963)  
A classification of Micrococci and Staphylococci based on physiological and Bio-chemical tests, *J. Gen. Microbiol.*, 30, 409-427 pp.
- BAIRD-PARKER, A. C. (1965),  
The classification of staphylococci and Micrococci from Worldwide Sources, *J. Gen. Microbiol.*, 38, 363 - 387 pp.
- BAIRD-PARKER, A. C. (1966)  
Methods for classifying Staphylococci and Micrococci, in 'Identifications methods for microbiologists, Part A' (B. M. Gibbs and F. A. Skinner Eds.), Academic Press, London, 59-64 pp.
- BLAKE I. (1935)  
Some observations on the bacterial flora of the alimentary tract of certain salmonidae, in 'Final. Report of the Furunculosis Committee', Scotland, Edinburgh : H. M. S. O., 60-67 pp.
- BRAMSNAES, F. (1965)  
Handling of fresh fish, in 'Fish as Food' Vol, 4, (G. Borgstrom Eds.), Academic Press, New York, London, 1-63 pp.
- BUCHANAN, R. E. and GIBBONS, N. E. (1974)-  
Bergey's Manual of Determinative Bacteriology, 8th Edn., Williams and Wilkins Co., Baltimore 217-632 pp.
- COWAN, S. T. (1977)  
Cowan and Steel's Manual for the identification of Medical Bacteria, 2nd Edn. (Revised by S. T. Cowan) Cambridge University Press, London.
- FRYER, G. and ILES, T. D. (1972)  
The cichlid fishes of the Great Lakes in Africa, their biology and evolution, Oliver and Boyd, Edinburgh.
- GIBBONS, N. E. and REED, G. B. (1930)  
The effect of autolysis in sterile tissues on subsequent bacterial decomposition *J. Bacteriol.*, 19, 73-88 pp.
- GORDEN, R. E. and Smith, N. R. (1949)  
Aerobic spore forming bacteria, *J. Bact.*, 58, 327-429 pp.
- HARRIGAN, W. F. and McCANCE, M. E. (1976)  
Laboratory methods in Food and Dairy Microbiology (Revised edition prepared by W. F. Harrigan), Academic Press London, New York.
- HENDRIE, M. S. and SHEWAN J. M. (1966)  
The identification of certain *Pseudomonas* species, in 'Identification methods for micro-biologists, Part A' (B. M. Gibbs and F. A. Skinner Eds.), Academic Press, London, New York, 1-7 pp.
- KATUGAMPOLA, D. S. and ASSIM, T. H. (1958)  
Coliform organisms in domestic water supplies in Ceylon, *Cey. Journ. of Med. Scie.*, 9, 95-101 pp.
- LISTON, J. (1956)  
Quantitative variations in the bacterial flora of flat fish, *J. Gen. Microbiol.*, 15, 305-314 pp.
- MARGOLIS, L. (1953)  
The effect of fasting on the bacterial flora of the intestines of fish, *J. Fish. Res. Board. Can.*, 10, 62-63 pp.
- MENDIS, A. S. and FERNANDO, C. H. (1962)  
A guide to the fresh-water fauna of Ceylon, *Bull. Fish. Res. Stn., Ceylon*, Vol. 12.



- SCHOLES, R. B. and SHEWAN, J. M. (1964)  
The present status of some aspects of marine microbiology, in 'Advances in marine biology' Vol. 2, (F. S., Russell Eds.) Academic Press, London, New York, 133-169 pp.
- SHEWAN, J.M. (1961)  
The microbiology of sea water fish, in 'Fish as Food, Vol. 1, (G. Borgstrom Eds.), Academic Press, New York London 487-560 pp.
- SHEWAN J. M. (1971)  
The microbiology of fish and fishery products, *J. Appl. Bact.*, **34** (2) 299-315 pp.
- SHEWAN, J.M.HOBBS, G. and (1967)  
The bacteriology of fish spoilage and preservation in 'Progress in industrial microbiology,' Vol. 6 (D.J.D, Hockenull Eds.), London Heywood Books, 169-208 pp.
- SHEWAN, J. M. HOBBS, G. and HODGKISS, W. (1960)  
A determinative scheme for the identification of certain genera of Gram-negative bacteria with special reference to Pseudomonadaceae *J. Appl Bact.*, **23**, 379-390 pp.
- SHEWAN, J.M. HOBBS, G. and HODGKISS, W. (1960)  
The *Pseudomonas* and *Achromobacter* groups of bacteria in the spoilage of marine white fish, *J. Appl. Bact.*, **23**. 463-468 pp.
- SMITH, N.R., GORDEN, R.E. and CLARK, F.E. (1952)  
Classification and identification scheme for Aerobic sporeforming bacteria, U.S. Dept. Agric. Monograph No. 16, Washington, D.C.
- SUB COMMITTEE: (1965)  
Minutes of 1st Meeting of Sub-Committee on taxonomy of Staphylococci and Micrococci *Inten. Bull. Bacteriol Nomenclature. Taxonomy*, **15**, 105-108 pp.
- TRUST, T. J. BULL, L.M. CURRIE B.R. and BUCKLEY, J.T. (1979)  
Obligate anaerobic bacteria in the gastrointestinal microflora of the Grass Carp (*Ctenopharyngodon idella*), Gold fish (*Carassius auratus*) and Rainbow Trout (*Salmo gairdner*), *J. Fish. Res. Board. Can.*, **36**, 1174-1179 pp.
- TRUST, T.J., MAC INNES, J.I. and BARTLETT, K.H. (1978)  
Variations in the intestinal microflora of Salmon fishes, in 'Microbiol Ecology' (M.W. Loutit and J.A.R.Miles Eds.), Springer-Verlag, Berlin, Heidelberg, New York, 250-254 pp.
- WOLF, J. and BARKER, A.N. (1968)  
The genus *Bacillus*-Aids to the identification of its species, in Identification Methods for Microbiologists, Part B', (B.M. Gibbs and D.A. Shapton Eds.), Academic Press, London, New York, 91-109 pp.
- WOOD, E.J.F. (1967)  
Microbiology of ocean and estuaries, Elsevier Oceanographic series, Vol. 3, Elsevier publishing company, Amsterdam, London, New York.